




AMERICAN JOURNAL OF ADVANCES IN NURSING RESEARCH

Journal homepage: www.mcmed.us/journal/ajanr



A STUDY TO ASSESS THE EFFECTIVENESS OF VIDEO- ASSISTED TEACHING CUM DEMONSTRATION ON ORAL HEALTH AMONG SCHOOL CHILDREN IN SELECTED SCHOOL AT KANYAKUMARI DISTRICT

K. Anilet Anandhy*

Professor, Department of child health nursing, Bhagyodaya Tirth Nursing College, Sagar, Madhya Pradesh, India.

Article Info

Received 25/01/2021

Revised 15/02/2021

Accepted 17/03/2021

Key word: Oral health, Video teaching cum demonstration, Effectiveness

ABSTRACT

A study to assess the effectiveness of video-assisted teaching cum demonstration on oral health among school children in selected school at kanyakumari district. The objectives of the study were to assess the knowledge regarding Oral Health among school children. To determine the effectiveness of Video Assisted Teaching cum demonstration among school children. To find out the association between the level of knowledge with their selected demographic variables through post-test. Pre experimental one group pre-test, post-test design was adopted for this study. The conceptual framework was based on Pender's modified Health Promotion Model. The study was conducted in Don Bosco matriculation school midalakadu, kanyakumari. Non probability purposive sampling technique was adopted to select the desired sample. The sample size was 60 children respectively. Structured interview questionnaire was used to assess the knowledge and practice regarding oral health among school children. The data collection tool was validated by experts and was found to be valid. The group was shown videos and demonstrated the steps of brushing and flossing. The result revealed that "t" value was 9.31, it was significant at $p < 0.05$ level. Hence the stated hypotheses H1 were accepted. It is inferred that video assisted teaching cum demonstration is effective in improving oral health knowledge and practice among school children.

INTRODUCTION

Children are assets of our country. Today's children are tomorrow's leaders. They form 38%-40% of our general population [1]. The children are one third of our population and all of our future. There are about 200.6 million children belonging to 6 to 12 years globally [2]. Children in school age are prone to get specific health problems [3].

Corresponding Author

K. Anilet Anandhy

Email: -aniletanandhy@gmail.com

The various acute and chronic conditions which can be encountered during school period anywhere in the world include oral health problems, Dental caries, Diarrhea, Worm Infestations, Hepatitis, Anemia, Scabies, Eczema, Acne, Influenza, Pneumonia, Diphtheria, Asthma, Fracture, Measles, Mumps, Chickenpox, Urinary Tract Infection, Meningitis, Tuberculosis, Eye and Ear Infections [4]. The school age child has multitude of problems, among them one of the most existing problem is related to oral health [5]. The goal of WHO "Health for



all by the year 2025” includes oral health also [6].

Oral health has also been found to profoundly influence the quality of life. Oral health is an integral component of general health [7]. Research in the past few years has revealed the causal link between oral diseases and systemic diseases [8]. Oral health is the royal way to overall health, wellbeing and quality of life [9]. It acts as a mirror that reflects general health/disease According to Osler “Oral cavity is a mirror of rest of the body “Oral health is essential for general health and well-being throughout the lifespan and is a mark of overall health status [10]. Research and other advances in oral health have led to safe and effective means of maintaining oral health and preventing dental caries and periodontal disease [11]. India, a developing country faces many challenges in rendering oral health needs. The majority of Indian population resides in rural areas, of which more than 40% constitute children [12]. These children cannot avail dental facilities due to inaccessibility, financial constraints and stagnation of public dental healthcare services [13]. This entails the health professional to adopt a more practical approach to achieve primary prevention of oral diseases [14]. The most viable solution seems to be dental and oral health education [15, 16].

Need for the study:

India is facing many challenges in rendering oral health care to the rural masses. Out of these 70-72% residing in rural areas more than 40% constitute children. This report is based on research survey with respect to different parameters i.e. Oral hygiene practices, dietary pattern, tobacco smoking & chewing, media habits and awareness regarding oral and dental treatment to get the complete overview of the existing oral health related problems and the factors responsible for poor oral health among rural children. WHO reported that 60-90% of school children worldwide have dental cavities and an oral health survey in India showed that the prevalence of oral and dental diseases is 90%.

Mehta A, Kaur G (2012) conducted a pre-experimental study to assess oral health related knowledge, attitude, and practices among 6- 12-year-old school children studying in rural areas of Panchkula, India. The ultimate goal was to implement an oral health-promotion program in this area .A total of 440 children 216 males and 224 females from 12 schools were included in this study. All the participants were requested to complete a 13-question closed-ended questionnaire. Only 25% of the participants said that they cleaned their teeth more than once in a day. 32% percent did not clean their teeth daily. Based upon the findings of the present study, the knowledge, attitude, and practices of the surveyed children with regard to oral health is poor. Hence, there is a need for regular oral health education of

the children, as well as their parents and school teachers.

Awareness regarding the importance of oral hygiene has significantly increased in the developed countries, but contrary to that, the modern dietary lifestyle habits are posing a greater risk for oral health. Healthy teeth not only enable you to look and feel good, they make it possible to eat and speak properly. Good oral health is important to your overall well-being.

There is a need of health education to all people. It has got preventive, promotive and rehabilitative dimensions. The school children can be an excellent mode to transmit information. Here the investigator hope that they can be an effective messengers of health, to other children, to their parents, to the family and finally to reach out the community. Hence the investigator felt that by doing this kind of study, will focus on prevailing problems of oral health and findings help to develop effective teaching program in order to modify the knowledge and practices of school children; So that the general health related complications among the future adults may be prevented leading to healthy generations.

MATERIALS AND METHODS:

A study was conducted in 2013 using pre-experimental research design (one group pre-test and post-test design) to assess the effectiveness of video assisted teaching cum demonstration on oral health among school children in selected school at Kanyakumari district. 60 schoolers were selected adopting non-probability purposive sampling technique. The tool has two parts, Part I: Demographic variables, Part II: Structured knowledge questionnaire to assess the knowledge regarding oral health among school children. Descriptive and inferential statistics.

RESULTS:

The study reveals that Regarding age 17(28.3%) belongs to age group of 7 years, 37(61.6%) belongs to age group of 8 years, 6(10%) belongs to age group of 9 years. Regarding Gender 26(43.3%) are males and 34(56.6%) are females. Regarding type of family 18(30%) belongs to joint family and 42(70%) belongs to nuclear family. Regarding number of siblings 20(33.3%) has no siblings, 18(30%) has one sibling, 22(36.6%) has more than one siblings. Regarding education of father 18(30%) were illiterates, 10(16.6%) were primary, 5(8.3%) were high school, 10(16.6%) were higher secondary, and 17(28.3%) were degree. Regarding education of mother 10(16.6%) were illiterates, 2(3.3%) were primary, 10(16.6%) were high school, 21(35%) were higher secondary and 17(28.3%) were degree. Regarding occupation father 18(30%) are coolies, 12(20%) are self employers, 20(33.3%) are private employers, 8(13.3%) are government employers and 2(3.3%) are



unemployed. Regarding occupation mother 7(11.6%) are coolies, 8(13.3%) are self employers, 7(11.6%) are private employers, 6(10%) are government employers and 32(53.3%) are home makers. Regarding monthly income of the family 3(5%) earns Rs<5000, 14(23.3%) earns 5001-10,000, 29(48.3%) earns 10,001 -20,000, 14(23.3%) earns >20,000. Regarding tooth brushing media 55(91.6%) were brush users and 5(8.3%) were non brush users. Regarding

previous knowledge about oral health 21(35%) has previous knowledge and 39(65%) has no previous knowledge on oral health. Regarding number of visits to dentist 28(46.6%) has never visited, 8(13.3%) visited once, 10(16.6%) visited twice, 14(23.3%) visited more than two times. number of visits to dentist in a year.

Table 1: Frequency and percentage distribution of demographic variables

SL. NO	DEMOGRAPHIC VARIABLES	FREQUENCY	PERCENTAGE %
1	Age years		
	7yrs	17	28.3
	8yrs	37	61.6
	9yrs	6	10
2	Sex		
	Male	26	43.3
	Female	34	56.6
3	Type of family		
	Joint	18	30
	Nuclear	42	70
4	No of siblings		
	None	20	33.3
	One	18	30
	More than one	22	36.6
5	Education of father		
	Illiterate	18	30
	Primary	10	16.6
	High school	5	8.3
	Higher secondary	10	16.6
	Degree/Equivalent	17	28.3
6	Education of mother		
	Illiterate	10	16.6
	Primary	2	3.3
	High school	10	16.6
	Higher secondary	21	35
	Degree/Equivalent	17	28.3
7	Occupation of father		
	Coolie	18	30
	Self employed	12	20
	Private employed	20	33.3
	Government employed	8	13.3
	Unemployed	2	3.3
8	Occupation of mother		
	Coolie	7	11.6
	Self employed	8	13.3
	Private employed	7	11.6
	Government employed/	6	10
	Home maker	32	53.3



9	Monthly income <5000	3	5
	5001-10,000	14	23.3
	10,000-20,000	29	48.3
	>20,000	14	23.3
10	Tooth brushing media Brush user	55	91.6
	Non brush user	5	8.3
11	Previous knowledge about oral health Yes	21	35
	No	39	65
12	No of visit to dentist in a year None		
	Once	28	46.6
	Twice	8	13.3
	More than 2 times	10	16.6
		14	23.3

The level of knowledge score is 9(15%) for inadequate knowledge, 10(17%) for moderately adequate knowledge and 41(68%) for adequate knowledge.

Table 2: Frequency and percentage distribution of knowledge after video assisted teaching cum demonstration.

Level of Knowledge	Classification of Respondents	
	Post test	
	Number	Percentage
Inadequate (<0-40% of score)	9	15
Moderate (41-60% of score)	10	17
Adequate (>61 % of score)	41	68
Total	60	100

It is inferred that there was significant association only between the level of knowledge of children with their age.

Table 3: Mean Standard Deviation and independent “t” test value of level of knowledge on oral health before and after video assisted teaching cum demonstration.n=60

S NO	Level of knowledge and practice on oral health	Mean	Standard deviation	“t” Test value
1.	Post Test	9.98	3.27	9.318*
2.	Pre Test	4.65	3.02	

* Significant at level $p < 0.05$

Table 2 shows that the mean value for the post test is 9.98 and the standard deviation is 3.27 and the mean pre-test value is 4.65 and standard deviation is 3.02. The tabulated “t” value is 1.77 and the obtained “t” value is 9.318, it is significant at $p < 0.05$ level. H_1 is accepted.



Table 4: Frequency, percentage and Chi-square distribution on the level of knowledge among school children with their selected demographic variables.

Demographic Variables	Inadequate		Moderate		Adequate		Chi-square
	n	%	n	%	n	%	
Age years							
7yrs	4	7	0	0	9	15	9.86
8yrs	4	7	6	10	27	45	p<0.05
9yrs	4	7	3	5	3	5	S
Gender							1.75
Male	4	7	6	10	15	25	P<0.05
Female	4	7	4	7	27	45	NS
Type of family							0.32
Joint	3	5	2	3	9	15	P<0.05
Nuclear	6	10	8	13	32	53	NS
Education of father							5.68
Illiterate	2	3	1	2	13	22	P<0.05
Primary	1	2	2	3	7	12	NS
High school	0	0	0	0	5	8	
Higher secondary	3	5	1	2	5	8	
Degree	1	2	3	5	16	27	
Education of mother							3.27
Illiterate	3	5	1	2	3	5	P<0.05
Primary	0	0	1	2	1	2	NS
High school	2	3	4	7	4	7	
Higher secondary	1	2	2	3	2	3	
Degree	3	5	9	15	24	40	
Occupation of father							2.47
Coolie	4	7	2	3	13	22	P<0.05
Self employed	2	3	3	5	7	12	NS
Private employed	2	3	4	7	6	10	
Government employed	2	3	3	3	3	3	
Unemployed	3	5	3	3	3	3	
Occupation of mother							3.4
Coolie	2	3	3	5	7	12	P<0.05
Self employed	2	3	4	7	11	18	NS
Private employed	2	3	3	5	5	8	
Government employed	3	5	4	7	6	10	
	3	5	3	5	31	52	



Home maker							
Monthly income	0	0	0	0	2	3	
<5000	3	5	4	7	13	22	0.85
5001-10,000	8	13	3	5	2	3	P<0.05
10,000-20,000	17	28	5	8	3	5	NS
>20,000							
Tooth brushing media							1.25
Brush user	1	2	0	0	4	7	P<0.05
Non brush user	8	13	10	17	37	62	NS
Previous knowledge							1.25
about oral health	5	8	0	0	4	7	P<0.05
Yes	8	13	10	17	33	55	NS
No							
No of visit to dentist in a year	4	7	5	8	19	32	1.25
None	1	2	1	2	6	10	P<0.05
Once	1	2	1	2	8	13	NS
Twice	1	2	3	5	10	17	
More than 2 times							

DISCUSSION AND CONCLUSION

Children are future of the country. Healthy children are important for healthy nation[17-20]. India remains one of the highest ranking in the world in terms

of number of children suffers with various disease in which oral problem is one among them. So it is important to create awareness among the pediatric population.

REFERENCES

1. Vangipuram, S., Jha, A., Raju, R., & Bashyam, M. (2016). Effectiveness of peer group and conventional method (Dentist) of oral health education programme among 12-15 year old school children - A randomized controlled trial. *Journal of Clinical and Diagnostic Research*. <https://doi.org/10.7860/JCDR/2016/17725.7844>.
2. Shilpa, pm, & Swamy, P. (2015). A study to evaluate the effectiveness of role play on knowledge regarding oral hygiene among higher primary school children in selected school at Tumkur. *IOSR Journal of Nursing and Health Science Ver. I*.
3. Alhayek, A. I. A., Alsulaiman, M. J., Almuhanha, H. A., Alsalem, M. A., Althaqib, M. A., Alyousef, A. A., Alabdali, J. N., Alqahtani, S. A., & Ansari, S. H. (2018). The effect of conventional oral health education versus animation on the perception of Saudi males in primary school children. *Journal of International Oral Health*. https://doi.org/10.4103/jioh.jioh_86_18.
4. Shettigar, S., Kamath, A., Alva, G. L., T., L., & Raju, N. J. (2013). TRAINING ON IMPROVING THE COMPETENCY LEVEL OF SELFADMINISTRATION OF INSULIN AMONG TYPE 2 DIABETES PATIENTS. *Journal of Health and Allied Sciences NU*. <https://doi.org/10.1055/s-0040-1703676>.
5. Al-Omiri, M. K., Al-Wahadni, A. M., & Saeed, K. N. (2006). Oral Health Attitudes, Knowledge, and Behavior Among School Children in North Jordan. *Journal of Dental Education*. <https://doi.org/10.1002/j.0022-0337.2006.70.2.tb04074.x>.
6. Bellows, L. L., Johnson, S. L., Davies, P. L., Anderson, J., Gavin, W. J., & Boles, R. E. (2013). The Colorado LEAP study: Rationale and design of a study to assess the short term longitudinal effectiveness of a preschool nutrition and physical activity program. *BMC Public Health*. <https://doi.org/10.1186/1471-2458-13-1146>.
7. Kuyken, W., Weare, K., Ukoumunne, O. C., Vicary, R., Motton, N., Burnett, R., Cullen, C., Hennelly, S., & Huppert, F. (2013). Effectiveness of the Mindfulness in Schools Programme: Non-randomised controlled feasibility study. *British Journal of Psychiatry*. <https://doi.org/10.1192/bjp.bp.113.126649>.
8. Adab, P., Pallan, M. J., Lancashire, E. R., Hemming, K., Frew, E., Griffin, T., Barrett, T., Bhopal, R., Cade, J. E., Daley, A., Deeks, J., Duda, J., Ekelund, U., Gill, P., McGee, E., Parry, J., Passmore, S., & Cheng, K. K. (2015). A cluster-randomised controlled trial to assess the effectiveness and cost-effectiveness of a childhood obesity prevention programme delivered through schools, targeting 6-7 year old children: The WAVES study protocol Disease



- epidemiology - Chronic. *BMC Public Health*. <https://doi.org/10.1186/s12889-015-1800-8>.
9. Tercedor, P., Villa-González, E., Ávila-García, M., Díaz-Piedra, C., Martínez-Baena, A., Soriano-Maldonado, A., Pérez-López, I. J., García-Rodríguez, I., Mandic, S., Palomares-Cuadros, J., Segura-Jiménez, V., & Huertas-Delgado, F. J. (2017). A school-based physical activity promotion intervention in children: Rationale and study protocol for the PREVIENE Project. *BMC Public Health*. <https://doi.org/10.1186/s12889-017-4788-4>.
 10. Prasai Dixit, L., Shakya, A., Shrestha, M., & Shrestha, A. (2013). Dental caries prevalence, oral health knowledge and practice among indigenous Chepang school children of Nepal. *BMC Oral Health*. <https://doi.org/10.1186/1472-6831-13-20>.
 11. Vander Ploeg, K. A., Maximova, K., McGavock, J., Davis, W., & Veugelers, P. (2014). Do school-based physical activity interventions increase or reduce inequalities in health? *Social Science and Medicine*. <https://doi.org/10.1016/j.socscimed.2014.04.032>.
 12. Takehara, K., Ganchimeg, T., Kikuchi, A., Gundegmaa, L., Altantsetseg, L., Aoki, A., Fukuie, T., Suwabe, K., Bat-Erdene, S., Mikami, M., Mori, R., & Soya, H. (2019). The effectiveness of exercise intervention for academic achievement, cognitive function, and physical health among children in Mongolia: A cluster RCT study protocol. *BMC Public Health*. <https://doi.org/10.1186/s12889-019-6986-8>.
 13. Sanadhya, Y. K., Thakkar, J. P., Divakar, D. D., Pareek, S., Rathore, K., Yousuf, A., Ganta, S., Sobti, G., Maniar, R., Asawa, K., Tak, M., & Kamate, S. (2014). Effectiveness of oral health education on knowledge, attitude, practices and oral hygiene status among 12–15-year-old schoolchildren of fishermen of Kutch district, Gujarat, India. *International Maritime Health*. <https://doi.org/10.5603/imh.2014.0022>.
 14. Damle, S., Damle, D., Jain, S., Patil, A., & Chopal, N. (2014). Effectiveness of supervised toothbrushing and oral health education in improving oral hygiene status and practices of urban and rural school children: A comparative study. *Journal of International Society of Preventive and Community Dentistry*. <https://doi.org/10.4103/2231-0762.142021>.
 15. Martínez-Vizcaino, V., Mota, J., Solera-Martínez, M., Notario-Pacheco, B., Arias-Palencia, N., García-Prieto, J. C., González-García, A., Álvarez-Bueno, C., & Sánchez-López, M. (2015). Rationale and methods of a randomised cross-over cluster trial to assess the effectiveness of MOVI-KIDS on preventing obesity in pre-schoolers. *BMC Public Health*. <https://doi.org/10.1186/s12889-015-1512-0>.
 16. Haque, S. E., Rahman, M., Itsuko, K., Mutahara, M., Kayako, S., Tsutsumi, A., Islam, M. J., & Mostofa, M. G. (2016). Effect of a school-based oral health education in preventing untreated dental caries and increasing knowledge, attitude, and practices among adolescents in Bangladesh. *BMC Oral Health*. <https://doi.org/10.1186/s12903-016-0202-3>.
 17. Xu, F., Ware, R. S., Leslie, E., Tse, L. A., Wang, Z., Li, J., & Wang, Y. (2015). Effectiveness of a randomized controlled lifestyle intervention to prevent obesity among Chinese primary school students: Click-obesity study. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0141421>.
 18. Cooper, A. M., O'Malley, L. A., Elison, S. N., Armstrong, R., Burnside, G., Adair, P., Dugdill, L., & Pine, C. (2013). Primary school-based behavioural interventions for preventing caries. In *Cochrane Database of Systematic Reviews*. <https://doi.org/10.1002/14651858.CD009378.pub2>.
 19. Salmon, J., Booth, M. L., Phongsavan, P., Murphy, N., & Timperio, A. (2007). Promoting physical activity participation among children and adolescents. In *Epidemiologic Reviews*. <https://doi.org/10.1093/epirev/mxm010>.
 20. Monse, B., Benzian, H., Naliponguit, E., Belizario, V., Schratz, A., & Van Palensteinhelderman, W. (2013). The Fit for School health outcome study - A longitudinal survey to assess health impacts of an integrated school health programme in the Philippines. *BMC Public Health*. <https://doi.org/10.1186/1471-2458-13-256>.

