



THE ROLE OF *GONGRONEMA LATIFOLIUM* IN ATTENUATION OF CHLOROQUINE INDUCED NEPHROTOXICITY AND HEPATOXICITY

*¹Nnodim Johnkennedy,¹ Uduji Helen Ifeoma and ²Udujih Godwin Obinna

¹Department of Medical Laboratory Science, Faculty of Health Science, Imo State University owerri, Imo State, Nigeria.

²Department of Public Health Technology, Federal University of Technology Owerri, Imo State Nigeria.

ABSTRACT

The attenuated effect of *Gongronemalatifolium* against chloroquine induced nephrotoxicity and hepatotoxicity was investigated in wistar rats. Twenty wistar rats were divided into 4 groups of 5 rats per group. Animal in group one served as control while animal in group 2, 3 and 4 received single oral administration of chloroquine (970mg/kg body weight). However experimental group 3 and 4 were additionally treated with 250mg/kg and 500mg/kg body weight of leaf extract of *Gongronemalatifolium* respectively. The treatment lasted for 14days. The results obtained showed that the serum urea and creatinine significantly increased in group 2 than in group 1, but significantly reduced in group 3 and 4 when compared with group 2 (P<0.05). On the other hand, serum bilirubin, AST, ALT, and ALP levels were significantly increased in group 2, when compared with the control. However, the level of serum bilirubin, AST, ALT, and ALP levels dropped marginally in group 3 and 4. This observation probably implies that the leaf extract of *Gongronemalatifolium* attenuated the nephrotoxicity and hepatotoxicity induced by chloroquine. Hence the use *Gongronemalatifolium* could be helpful to patients with liver and kidney derangement.

Key words: Chloroquine, Nephrotoxicity, Hepatotoxicity, *Gongronemalatifolium*.

Corresponding Author

NnodimJohnkennedy

Email: -johnkennedy23@yahoo.com

Article Info

Received 12/03/2014; Revised 20/03/2014

Accepted 02/04/2014

INTRODUCTION

Chloroquine is a member of an important series of chemically related anti malarial agents, the quinolone derivatives. Being a 4-aminoquinoline, it is a rapidly acting blood schizontocide with some gametocytocidal activity [1]. It is a synthetic drug for effective treatment of malaria and the anti malarial drug of choice for many years in most parts of the world. In malaria endemic regions like south eastern part of Nigeria, where *Plasmodium falciparum* is sensitive to chloroquine, the drug is still useful for the treatment and prophylaxis of the infection, in spite of increasing prevalence of resistance of the parasite to the drug [2,3]. However, chloroquine treatment is often accompanied by serious side effect such as headache and visual disturbances. It has been reported to cause hepatic damage [4]. The free radicals have been implicated in many diseases including nephrotoxicity and

hepatotoxicity [5]. Experimental, clinical and epidemiological studies have provided evidence in support of the role of free radicals in the etiology of hepatotoxicity by anti malaria drugs [6,7]. When produced in excess, free radicals cause tissue injury. However, the natural antioxidants defense mechanisms can be inadequate and therefore some herbal plant can be of great importance. One of such plants is *Gongronemalatifolium* this plant *Gongronemalatifolium* is locally called “utazi by Igbos, “arokeke” by Yorubas, and Utasi” by the Efiks and Ibiobios in Nigeria. *Gongronemalatifolium* belongs to the family of Asclepiadaceae. It is an edible rainforest plant native to the South East part of Nigeria and has widely used in folk medicine [8]. It is a herbaceous shrub with yellow flowers and the stem that yields characteristic milky exudates when cut. Some phytochemicals such as β – sosterol,



lupenyl esters, Pregnancy ester, glucosides, essential oils and saponins are associated with parts of this herb [9]. *Gongronemalatifolium* which has been used in traditional medicine for treating diabetes, malaria, hypertension and as laxative. Also used as a spice and vegetable [10,11]. Since this plant has been described to be beneficial, the present study investigates the protective attenuation role of *G latifolium* in chloroquine induced nephrotoxicity and hepatotoxicity as there have been relatively few studies on this model.

MATERIALS AND METHODS

Plant materials

The *Gongronemalatifolium* leaves were obtained from Ekeonunwa market in Owerri Nigeria. The botanical identification and authentication was confirmed by Dr. C. Okere (Head of Department of Plant Science and Biotechnology, Imo State University, Owerri). The plant material was sun dried for seven days. The dried leaves of *Gongronemalatifolium* were milled to get a fine powder. The appropriate concentrations of the extract were made in distilled water for the experiment. Hence, the following concentrations: 250mg and 500mg were prepared.

Drugs

Chloroquine (Emzor) was purchased from a standard pharmacy shop in Owerri, Imo State Nigeria. The tablets were dissolved in distilled water according to the required concentrations required for administration to the Wistar rats on the basis of their body weight.

Experimental Animals

The Wistar albino rats weighing between 170 and 220g, ages (8-10 weeks) were used in the study. These animals were obtained from the Animal House of College of Medicine and Health Sciences, Imo State University, Owerri Nigeria. They were kept under standard laboratory conditions, fed with commercial growers mash, product of Tops Feeds Ltd, Sapele, Nigeria. Water and feed were provided *ad libitum*. The animals were left for two weeks to acclimatize and then divided into groups for experimentation.

Experimental Design

The animals were randomly assigned to four experimental groups (n = 5 x 4group). The first group of animals which served as control was given distilled water. Group II, III and IV were given chloroquine (970mg/kg body weights), chloroquine and *G latifolium* extract (250mg/kg body weight), chloroquine and *G latifolium* extract (500mg/ kg body weight) respectively for 14 days. In all groups the drug was administered through oral route using a feeding tube attached to a 5ml syringe. All animals were allowed free access to food and water throughout the experiment.

Blood Collection

Twenty four hours after the last doses were administered, the animals were anaesthetized with chloroform vapor, quickly brought out of the jar and sacrificed. Whole blood was collected by cardiac puncture from each animal into clean dry test tubes. The blood was allowed to stand for about 15 minutes to clot and further spun in a Westerfuge centrifuge (model1384) at 10,000g for 5 minutes. Serum was separated from the clot with Pasteur pipette into sterile sample tubes for the measurement of the biochemical parameters.

Biochemical Analysis

Serum urea and creatinine were measured using standard method[12]. Serum AST and ALT were assayed by the method of Reitman and Frankel(1957). ALP was determined by the method of king and king (1954) [13]. Also, serum bilirubin was determined by the method of Jendrassik and Groff (1938) [14].

Statistical analysis

The results were expressed as mean ± standard deviation. The statistical evaluation of data was performed by using students t- test. The level of significance was calculated at P<0.05.

The results were expressed as mean ± standard deviation. The statistical evaluation of data was performed by using one-way ANOVA (Analysis of variance) followed by Duncan’s multiple range test [15].

RESULTS

Table 1. Effect of *G. latifolium* on chloroquine induced nephrotoxicity Group Treatment urea (mg/dl) creatinine(mg/dl)

1	Control	19.2 ±4.1	0.6 ±0.02
2	Chloroquine	39.97 ±4.7 *	1.8 ±0.03*
3	Chloroquine+G.L250mg	27.43±6.2*	1.41±0.03*
4	Chloroquine+G.L500mg	22.59±6.0*	1.10±0.02*

*Significantly different from control (P<0.05)

Table 2. Effect of *G. latifolium* on chloroquine induced hepatotoxicity Group Treatment AST(iu/l) ALT(iu/l) ALP(iu/l) Bilirubin(mg/dl)

1	Control	14.2 ±2.1	13.11±2.8	66.22±9.020.6± 0.02
---	---------	-----------	-----------	---------------------



2	Chloroquine	33.27±3.1*	24.31± 2.9	93.8 ±8.73*1.9 ±0.02*
3	Chloroquine+G.L250mg	24.93±3.6*	21.42 ± 3.4*	85.71±8.03*1.42±0.04 *
4	Chloroquine+G.L500mg	20.37±3.0*	17.51± 3.0	79.43±6.10* 1.20±0.04*

*Significantly different from control (P<0.05)

G.L *Gongronemalatifolium*

DISCUSSION

The commonest biochemical parameters regarded as indicators of liver damages are AST, ALT, ALP and bilirubin while urea and creatinine is important renal indicator for kidney damage. The damage to the hepatocellular cells and renal cells resulted in the increase in these biochemical parameters [16,17]. In this study, chloroquine administration in the dose of 970mg/kg body weight of rats results in increase serum bilirubin, AST, ALT, ALP and activities as well as urea and creatinine. The increase is roughly proportional to the extent of liver enzymes in serum following anti malaria administration as has been earlier reported [15]. The liver and renal cell damage may be associated with the generation of reactive oxygen species (ROS) by chloroquine overdose which are also partly responsible for their anti-malarial effects, hence the harmful effects were considered to be caused by ROS produced during peroxide formation [18, 19]. The level of hydroxyl and peroxide radical induced by

chloroquine treatment may be responsible for the hepatic and renal impairment in Wistar rats [20]. The observed increased in enzyme activities may be probably due to a leakage of cytoplasmic enzyme into circulation as a result of inflammation of the liver cells. However, the simultaneous administration of *G. latifolium* along with CQ significantly reduced the effect of chloroquine by lowering bilirubin, AST, ALT and ALP activities as well as urea and creatinine. In the same vein, significantly increased serum bilirubin concentration in group 2 was significantly reduced on the administration of *G latifolium* when compared. This result suggests that *G latifolium* offer attenuation effect by preserving the structural integrity of renal and hepatocellular membrane against chloroquine.

In conclusion the present investigation indicates that *Gongronemalatifolium* exerted significant protection against chloroquine induced nephrotoxicity and hepatotoxicity

REFERENCES

1. Ekanem DJ, Weisfeld JS, and Salako LA (1990). Sensitivity of *Plasmodium falciparum* to chloroquine and sulphadoxine in Nigerian children. *Bulletin of the World Health Organization*, 68, 45-52.
2. Sowunmi A, Fehintola FA, Falade AG, Akinyinka OO, Oduola AMJ. (2000). Comparative efficacy of chloroquine plus chlorpheniramine alone or in sequential combination with sulfadoxine-pyrimethamine for treatment of acute uncomplicated falciparum malaria in Nigerian children. *Ann. Trop. Med. Para*, 94 (3), 209- 217.
3. Pari I and Amali RD. (2005). Protective role of tetrahydrocurmin (THC) an active principle of turmeric on chloroquine induced hepatotoxicity in rats. *J. Pharm PharmaceutSci*, 8(1), 115-123.
4. Pari I and Murugavel P. (2004). Protective effect of a lipoic acid against CQ induced hepatotoxicity in rats. *J APP Toxicol* ; 24 : 21-26 Reitman, S, Frankel S. Transaminases (1957). *Am J ClinPathol*, 28, 56.
5. Nnodim JK, Emejulu A, Amaechi A and Nwosunjoku EC. (2010). Alterations in biochemical parameters of wistar rats administrated with sulfadoxine and pyrimethamine (*Fansidar^R*). *Almeen J Med Sci*, 3 (4), 317-321.
6. Ali Y, Munir O and Vahit B. (2001). The antioxidant activities of leaves of *Cydonia vulgaris*. *Tur J Med Sci.*, 31, 23-2710.
7. Nnodim Jk, Afolabi EA, Udujih HI, Okorie H, Nwobodo EI and Nwadike CN. (2012a). Alterations in some biochemical indices of hepatic function in tuberculosis patients on antituberculosis therapy. *Indian Journal of Medicine and Healthcare*, 1(1), 12-15.
8. Nnodim JK, Emejulu A, Ihim A and Udujih HI. (2012b). Influence of *Gongronemalatifolium* on some biochemical parameters in alloxan induced diabetes. *International Journal of Analytical Pharmaceutical and Biomedical Sciences*, 1(1), 13-17.
9. Edet EE, Akpanabiatu MI, ENo AE, Umoh IB, and Itam EH. (2009). Effect of *GongronemaLatifolium* Crude leaf extract on some cardiac enzymes of alloxan induced diabetic rats. *African J Biochem Res*, 3(11), 366 – 369.
10. Morebise O, Fafunso MA, Makinde JM, Olayide OA, and Awe E. (2002). Anti inflammatory property of *Gongronemalatifolium*. *Phytother Res*, 16, 575 – 577.
11. Ugochukwu NH, Babady NE, Cobourne M, and Gasset SR. (2003). The effect of *Gongronemalatifolium* extracts on serum lipid profile and oxidative stress in hepatocytes of diabetic rats. *J. Bioscience*, 28(1), 1-5.
12. Cheesbrough, M. (2003). *District Laboratory practice in tropical countries part 1* Cambridge University Press 340-348.
13. King E, Jand King PR. (1954). Estimation of plasma phosphates by determination of hydrolyzed phenol with amino antipyrine. *J Chem Path*, 7, 322 – 32.
14. Jendrassik L and Groff P. (1938). Estimation of Bilirubin. *Biochem*, 297, 81.



15. Nwanjo H, U and Oze G. (2009). Acute hepatotoxicity following administration of artesunate in guinea pigs. *Internal J Toxicol*, 4, 1.
16. Nwanjo AU, Okolie NJ, Oze G, Okafor, MC, Nwosu D, Ajero, C and Anyaehie B. (2007). Halofantrine (antimalaria) toxicity in wistar rats. Biochemical evaluation of hepatic dysfunction. *Res J Med Sci*, 102-105.
17. Nnodim, JK, Ihim A and Uduji HI. (2012c). Attenuation of chloroquine-induced hepatotoxicity and renal damage by *Gnetumbucholzianum* leaf extract. *New Zealand Journal of medical Laboratory Science*, 66, 46-47.
18. Valko M, Leibfritz D, Monocol J, Cronin, MT, Mazur M and Tester J. (2007). Free radicals and anti oxidants in normal physiological functions and human disease. *Int. J. Biochem Cell Biol*, 39, 44-84.
19. Nnodim, Jk, Ujowundu CO, Udujih HI, Okorie H, Nwobodo EI and Nwadike CN. (2012d). Hepatocellular enzyme activities and protein level following administration of Vitamins during chloroquine induced hepatotoxicity in wistar rats. *Indian Journal of Innovation and development*, 1(4), 285-288.
20. Iniaghe MO, Malomo SO and Adebayo JO. (2008.) Hepatoprotective effect of the aqueous extract of leaves of *Acalypharacemosa* in carbon tetrachloride treated rats. *J Medicinal Plant Research*, 2 (10), 301-305.
21. Ita SO, Etim OE, Ben EE and Ekpo OF. (2007). Haemotopoietic properties of ethanolic leaf extract of *Ageratum conyzoides* (goat weed) in albino rats. *NIG J PhysiologSci*, 22, 83-87.
22. Tietz NW. (1995). Clinical guide to laboratory tests Third edition, WB Saunders Company Philadelphia, 518-519.

