



A CASE REPORTING OSMF IN 40 YEAR OLD PATIENT

*¹Gaurav Solanki,¹Namita Lohra,¹Jaya Lohra,²Renu Solanki

¹Jodhpur National University, Jodhpur, Rajasthan, India.

²Lachoo Memorial College of Science and Technology, Rajasthan, India.

Corresponding Author

GauravSolanki
Email:-drguravsolanki@yahoo.com

Article Info

Received 12/02/2014; Revised 16/02/2014
Accepted 30/03/2014

ABSTRACT

Oral submucous fibrosis a chronic disease of oral mucosa characterized by inflammation and progressive fibrosis followed by stiffening of healthy mucosa resulting in difficulty in opening the mouth. The treatment of patients with OSMF depends on the degree of clinical involvement. If the disease is detected at a very early stage, cessation of the habit is sufficient but at later stages, the treatment is relatively difficult. The present article focuses on a patient suffering from OSMF and also discusses about its signs and symptoms.

Key Words:OSMF, Etiology, Lab Findings etc.

INTRODUCTION

Oral submucous fibrosis a chronic disease of oral mucosa characterized by inflammation and progressive fibrosis followed by stiffening of healthy mucosa resulting in difficulty in opening the mouth [1]. It is seen today that areca nut quid plays a major role in the etiology. The younger generation is very much addicted to these products especially gutkha and panmasala. The condition is well recognized for its malignant potential. The pathogenesis of the disease is not well established, but the cause of OSMF is believed to be multifactorial. Factors include areca nut chewing, ingestion of chilies, genetic and immunologic processes, nutritional deficiencies and other factors [2]. Iron deficiency anemia, vitamin B complex deficiency, and malnutrition are promoting factors that derange the repair of the inflamed oral mucosa, leading to defective healing and resultant scarring. The rate varies from 0.2-2.3% in males and 1.2-4.57% in females in Indian communities [3].

CASE REPORT

A 40-year-old man presented with 1 year history of burning sensation in the oral mucosa with chillies and difficulty in opening the mouth on right side since 6 months. No history of any mucosalerosion or ulcers was

present and the medical history was non significant. History of chewing tobacco with lime and arecanut was there for the last 5 years. He used to keep betel leaf with slaked lime and arecanut in the form of paan on right side 7-8 times a day. On examination, the right side oral mucosa was white and shiny. On right side, against the buccal aspect of molar teeth there was a firm band extending from the molar area upto the angle of mouth. The opening of the mouth was reduced. All the routine investigations were normal. Biopsy from the right buccal mucosa showed the presence of OSMF.



DISCUSSION

OSMF is widely prevalent in all age groups and across all socioeconomic strata in India. Generally, patient age ranges from 11-60 years most patients are aged 45-54 years and chew betel nuts 5 times per day. OSMF has a high rate of morbidity because it causes a progressive inability to open the mouth, resulting in eating and consequent nutritional deficiencies [4]. OSMF also has a significant mortality rate because it can transform into oral cancer, particularly squamous cell carcinoma, at a rate of 7.6%. Symptoms of OSMF include: Progressive inability to open the mouth (trismus) due to oral fibrosis and scarring, Oral pain and a burning sensation upon consumption of spicy foodstuffs, Increased salivation, Change of gustatory sensation, Hearing loss due to stenosis of the eustachian tubes, Dryness of the mouth, Nasal tonality to the voice, Dysphagia to solids (if the esophagus is involved), Impaired mouth movements etc [5]. Laboratory findings includes: Decreased hemoglobin levels, Decreased iron levels, Decreased protein levels, Increased erythrocyte sedimentation rate, Decreased vitamin B complex levels etc [6]. The treatment of patients with OSMF depends on the degree of clinical involvement. If the disease is detected at a very early stage, cessation of the habit is sufficient. Most patients

with OSMF present with moderate-to-severe disease. Moderate-to-severe OSMF is irreversible. Medical treatment is symptomatic and aimed at improving mouth movements. In patients with moderate OSMF, weekly sub mucosal intraregional injections or topical application of steroids may help prevent further damage [7]. The rationale for using placental extract (PE) in patients with OSMF derives from its proposed anti-inflammatory effect hence, preventing or inhibiting mucosal damage. The use of topical hyaluronidase has been shown to improve symptoms more quickly than steroids alone. The combination of steroids and topical hyaluronidase shows better long-term results [8]. Cox in 1996 said that dietary focus should be on reducing exposure to the risk factors, especially the use of betel quid and correcting any nutritional deficiencies, such as iron and vitamin B complex deficiencies [9, 10].

CONCLUSION

OSMF also has a significant high mortality rate and is widely prevalent in all age groups and across all socioeconomic strata in India. Patient education and public awareness regarding OSMF can be done to improve the condition and prevent future occurrence of OSMF.

REFERENCES

1. Pindborg J. (1980). Incidence and early forms of oral submucous fibrosis. *Oral surg. oral med. Oral pathol*, 50, 40-44.
2. Seedat A. (1988). Betel-nut chewing and submucous fibrosis in Durban. *S. Afr. med. j*, 74, 568-571.
3. Pindborg J. (1989). Oral submucous fibrosis, a review. *Ann. Acad. Med. Sing*, 18, 603-607.
4. Bhonsle B. (1987). Regional variations in oral submucous fibrosis in India. *Community dent. Oral epidemiol*, 15, 225-229.
5. Shiau Y. (1979). Submucous fibrosis in Taiwan. *Oral surg. oral med. oral pathol*, 47, 453-457.
6. Pindborg J. (1984). Pilot survey of oral mucosa in areca (betel) nut chewers on Hainan Island of the People's Republic of China. *Community dent. Oral Epidemiol*, 12, 195-196.
7. Ramanathan K. (1981). Oral submucous fibrosis, an alternative hypothesis as to its causes. *Med. j. Malaysia*, 36, 243-245.
8. Paissat K. (1981). Oral submucous fibrosis. *Int. J. Oral Surg*, 10, 307-312.
9. Lay M. (1982). Epidemiologic study of 600 villagers of oral precancerous lesions in Bilugyun, preliminary report. *Community Dent. Oral Epidemiol*, 10, 152-155.
10. Shear M. (1967). Oral submucous fibrosis, a possible case in a person of Caucasian descent. *Br. Dent. J*, 122, 343-346.

